



The Flu: What To Do If You Get Sick

On This Page

- [How do I know if I have the flu?](#)
- [What should I do if I get sick?](#)
- [What are the emergency warning signs?](#)
- [Do I need to go the emergency room if I am only a little sick?](#)
- [Are there medicines to treat the flu?](#)
- [How long should I stay home if I'm sick?](#)
- [What should I do while I'm sick?](#)

How do I know if I have the flu?

You may have the flu if you have some or all of these symptoms:

- fever*
- cough
- sore throat
- runny or stuffy nose
- body aches
- headache
- chills
- fatigue
- sometimes diarrhea and vomiting

*It's important to note that not everyone with flu will have a fever.

What should I do if I get sick?

Most people with the flu have mild illness and do not need medical care or antiviral drugs. If you get sick with flu symptoms, in most cases, you should stay home and avoid contact with other people except to get medical care.

If, however, you have symptoms of flu and are very sick or worried about your illness, contact your health care provider (doctor, physician's assistant, etc.).

Certain people are at greater risk of serious flu-related complications (including young children, elderly persons, pregnant women and people with certain long-term medical conditions) and this is true both for seasonal flu and novel flu virus infections. (For a full list of people at higher risk of flu-related complications, see [People at High Risk of Developing Flu-Related Complications](#)). If you are in a high risk group and develop flu symptoms, it's best for you to contact your doctor. Remind them about your high risk status for flu.

Health care providers will determine whether influenza testing and possible treatment are needed. Your doctor may prescribe antiviral drugs that can treat the flu. These drugs work better for treatment the sooner they are started.

What are the emergency warning signs?

In children

- Fast breathing or trouble breathing
- Bluish skin color
- Not drinking enough fluids
- Not waking up or not interacting
- Being so irritable that the child does not want to be held
- Flu-like symptoms improve but then return with fever and worse cough
- Fever with a rash

In addition to the signs above, get medical help right away for any infant who has any of these signs:

- Being unable to eat
- Has trouble breathing
- Has no tears when crying
- Significantly fewer wet diapers than normal

In adults

- Difficulty breathing or shortness of breath
- Pain or pressure in the chest or abdomen
- Sudden dizziness
- Confusion

- Severe or persistent vomiting
- Flu-like symptoms that improve but then return with fever and worse cough

Do I need to go the emergency room if I am only a little sick?

No. The emergency room should be used for people who are very sick. You should not go to the emergency room if you are only mildly ill. If you have the emergency warning signs of flu sickness, you should go to the emergency room. If you get sick with flu symptoms and are at high risk of flu complications or you are concerned about your illness, call your health care provider for advice. If you go to the emergency room and you are not sick with the flu, you may catch it from people who do have it

Are there medicines to treat the flu?

Yes. There are drugs your doctor may prescribe for treating the flu called “antivirals.” These drugs can make you better faster and may also prevent serious complications. See [Treatment - Antiviral Drugs](#) for more information.

How long should I stay home if I’m sick?

CDC recommends that you stay home for at least 24 hours after your fever is gone except to get medical care or for other things you have to do and no one else can do for you. (Your fever should be gone without the use of a fever-reducing medicine, such as Tylenol®.) You should stay home from work, school, travel, shopping, social events, and public gatherings.

What should I do while I’m sick?

Stay away from others as much as possible to keep from making them sick. If you must leave home, for example to get medical care, wear a facemask if you have one, or cover coughs and sneezes with a tissue. And wash your hands often to keep from spreading flu to others. Visit the [Taking Care of a Sick Person in Your Home](#) guide for more information.

View page in:

- [Español](#)



- Page last reviewed: January 26, 2012
- Page last updated: January 26, 2012
- Content source: [Centers for Disease Control and Prevention, National Center for Immunization and Respiratory Diseases \(NCIRD\)](#)



*Centers for Disease Control and Prevention 1600 Clifton Rd. Atlanta, GA 30333, USA
800-CDC-INFO (800-232-4636) TTY: (888) 232-6348 - [Contact CDC-INFO](#)*